New Patient Information

Patient Name:			DOB:		Age:
(Nombre de Paciente)			(Fed	cha de Nacimiento)	(Edad)
Address:		City:		ST:	Zip
(Direccion)		_	(Ciudad)	(Estado)	(Codigo)
SSN:	Home Phone#		Work:		Cell:
(Numero de seguro social)	Home Phone#(Telefono de casa)		(Telefon	o de Trabajo)	(Telefono de cellular)
Employer:			Phone#:		
(Nombre de Empleo)			(Telefono d	e Empleo)	
Spouse's Name:	*********	******	_ DOB:		Age:
(Nombre de Esposo/a)			(Fe	cha de Nacimento)	(Edad)
Spouse's Employer:			SSN:		
Spouse's Employer: (Empleo de Esposo/a)			(Nun	iero de seguro soci	al)
	Y 1	r e 4	•		
	Insurance]	(ntormat)	ion		
Primary Insurance:	DIC Mort	Lers Co	mo Phone#:		
(Nombre de Aseguransa Prima	(Informacion sobjection)			(Numero de	Telefono) .
Claims Address:					
(Direccion)		_	(Ciudad	ST: (Estado	(Codigo)
Insured's Name:		Group#_			
(Nombre de el persona que tien	ies el Aseguransa)	0.04.pn_	(1	Numero de Grupo)	
Marihar/ID#		► Doliavi#•	•		
Member/ID#:(Numero de miembro o ID)		Policy#:		Numero de Poliza)	
(Numero de intembro o 10)	Secondary I	ngurance	•	ivumero de 3 onza)	
	(Informacion sobre su Ase				
Secondary Insurance:	•		Phone#		
(Nombre de Aseguransa)				(Numero de	e Telefono)
Claims Address:		City:		ST:	_ Zip:
(Direccion)			(Ciudad)	(Estado	(Codigo)
Insured's Name:		Group#_			
(Nombre de el persona que tiene	es el Aseguransa)	Group#_	(N	umero de Grupo)	
lane u		70 le //	•		
Member/ID#:(Numero de miembro o ID)		Policy#: _		(Numero de Poliza	
(Numero de miembro o 12)		******	********	**********	'/ ************
Attorney's Name:			Phone#•		
Nombre de Abogado)			1 11011011.	(Telefono de s	u Abogado)
4 7 7		C:4		ст.	7:
Address Direction)		_ City:	(Ciudad)	ST: (Estado)	_Zip: (Codigo)
Date of Injury:	Who may we thank fo	er referring	von to our off	0e?	
Fecha que fue lastimado)	(Aquien le debemos las gr				
		-	•		•
'Patient's S ^{ig} nature: (Firma de Paciente)			Date:	(Fecha)	
\				,,, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	

AUTHORIZATION FOR THE RELEASE OF PROTECTED HEALTH INFORMATION (PHI) By Joseph J. Schifini, MD, LTD. (JJS)

Patient Name:			
	Last First M.I. (Previous or Other Name	es Used)	.
Address:			
Addiess.			
Date of Birth:			
If this Authoriza purpose below:	tion is for any purpose other than	the release of PHI for pe	rsonal reasons, please state the
	lease of medical records from:	Joseph J. Schifini, M 8925 W. Russell Roa Las Vegas, NV 891	ad Suite 200
5 1 1		-	
Please release red medical records to		Name:	
medical records (5.		
	•	Address: City:	
	•		
•		210	
		Telephone Number: (
		Fax Number: () -
I specifically auth	orize the use and disclosure of th	e following PHI: (Please	e provide a detailed
	e particular data and period of		g)
☐ Emergency Rec ☐ Clinic Records		spital Records liology Reports	
□ Lab Reports		liology Films	
☐ Shot Records	□ Patl	nology Reports	
□ Slides	□ Oth	er	<u>—</u>
This authorization	n will expire on the 180th day of th	ne signing unless a lesser	date is specified below:
information (PHI) as a information relating to treatment for drug or a notes maintained by a authorization at any tin 8925 W. Russell Roac revocation will not have revocation. If neither f disclosed pursuant to t laws. This Authorization	rization Form. I understand that I am giving lescribed above. The information to be used: (1) Acquired immunodeficiency syndrogalcohol abuse, or (3) mental or behavioral mental health provider, a separate authoring by notifying JJS in writing to Joseph J. Suite 200, Las Vegas, Nevada 89148 of we any effect on any information already usederal nor Nevada privacy law apply to the his authorization may be re-disclosed by the ton is voluntary and I may refuse to sign the exchange for the patient receiving treatments.	ed or disclosed pursuant to this me (AIDS) or (2) human immut health or psychiatric care. If y zation form must be completed. Schifini, MD, LTD Health In my intent to revoke this authoused or disclosed by JJS before the recipient of the information, the recipient and no longer promis Authorization Form. I unde	s authorization form may include unodeficiency virus (HIV) infection, you are requesting psychotherapy sessiod. I understand that I may revoke this formation Management Department, vization. I understand that such a US received my written notice of I understand that the information tected by federal or Nevada privacy
Signature of Pati	ent or Authorized Personal Rep	presentative	Date
Relationship to the	ne Patient (If signed by a Personal Re	presentative)	Date

HIPAA Notice of Privacy Practices

This notice was published and becomes effective on/or before April 14, 2003.

THIS NOTICE DESCRIBES HOW MEDICAL INFORMATION ABOUT YOU MAY BE USED AND DISCLOSED AND HOW YOU CAN GET ACCESS TO THIS INFORMATION. PLEASE REVIEW IT CAREFULLY.

This Notice of Privacy Practices describes how we may use and disclose your protected health information (PHI) to carry out treatment, payment or health care operations (TPO) and for other purposes that are permitted or required by law. It also describes your rights to access and control your protected health information. "Protected health information" is information about you, including demographic information, that may identify you and that relates to your past, present or future physical or mental health or condition and related health care services.

1. Uses and Disclosures of Protected Health Information

Uses and Disclosures of Protected Health Information

Your protected health information may be used and disclosed by your physician, our office staff and others outside of our office that are involved in your care and treatment for the purpose of providing health care services to you, to pay your health care bills, to support the operation of the physician's practice, and any other use required by law.

<u>Treatment</u>: We will use and disclose your protected health information to provide, coordinate, or manage your health care and any related services. This includes the coordination or management of your health care with a third party. For example, we would disclose your protected health information, as necessary, to a home health agency that provides care to you. For example, your protected health information may be provided to a physician to whom you have been referred to ensure that the physician has the necessary information to diagnose or treat you.

<u>Payment</u>: Your protected health information will be used, as needed, to obtain payment for your health care services. For example, obtaining approval for a hospital stay may require that your relevant protected health information be disclosed to the health plan to obtain approval for the hospital admission.

Healthcare Operations: We may use or disclose, as-needed, your protected health information in order to support the business activities of your physician's practice. These activities include, but are not limited to. quality assessment activities, employee review activities, training of medical students, licensing, and conducting or arranging for other business activities. For example, we may disclose your protected health information to medical school students that see patients at our office. In addition, we may use a sign-in sheet at the registration desk where you will be asked to sign your name and indicate your physician. We may also call you by name in the waiting room when your physician is ready to see you. We may use or disclose your protected health information, as necessary, to contact you to remind you of your appointment. We may use or disclose your protected health information in the following situations without your authorization. These situations include: as Required By Law, Public Health issues as required by law, Communicable Diseases: Health Oversight: Abuse or Neglect: Food and Drug Administration requirements: Legal Proceedings: Law Enforcement: Coroners, Funeral Directors, and Organ Donation: Research: Criminal Activity: Military Activity and National Security: Workers' Compensation: Inmates: Required Uses and Disclosures: Under the law, we must make disclosures to you and when required by the Secretary of the Department of Health and Human Services to investigate or determine our compliance with the requirements of Section 164.500.

Other Permitted and Required Uses and Disclosures Will Be Made Only With Your Consent, Authorization or Opportunity to Object unless required by law.

You may revoke this authorization, at any time, in writing, except to the extent that your physician or the physician's practice has taken an action in reliance on the use or disclosure indicated in the authorization.

Your Rights

The following is a statement of your rights with respect to your protected health information.

You have the right to inspect and copy your protected health information. Under federal law, however, you may not inspect or copy the following records; psychotherapy notes; information compiled in reasonable anticipation of, or use in, a civil, criminal, or administrative action or proceeding, and protected health information that is subject to law that prohibits access to protected health information.

You have the right to request a restriction of your protected health information. This means you may ask us not to use or disclose any part of your protected health information for the purposes of treatment, payment or healthcare operations. You may also request that any part of your protected health information not be disclosed to family members or friends who may be involved in your care or for notification purposes as described in this Notice of Privacy Practices. Your request must state the specific restriction requested and to whom you want the restriction to apply. Your physician is not required to agree to a restriction that you may request. If physician believes it is in your best interest to permit use and disclosure of your protected health information, your protected health information will not be restricted. You then have the right to use another Healthcare Professional.

You have the right to request to receive confidential communications from us by alternative means or at an alternative location. You have the right to obtain a paper copy of this notice from us, upon request, even if you have agreed to accept this notice alternatively i.e. electronically.

You may have the right to have your physician amend your protected health information. If we deny your request for amendment, you have the right to file a statement of disagreement with us and we may prepare a rebuttal to your statement and will provide you with a copy of any such rebuttal.

You have the right to receive an accounting of certain disclosures we have made, if any, of your protected health information.

We reserve the right to change the terms of this notice and will inform you by mail of any changes. You then have the right to object or withdraw as provided in this notice.

Complaints

You may complain to us or to the Secretary of Health and Human Services if you believe your privacy rights have been violated by us. You may file a complaint with us by notifying our privacy contact of your complaint. We will not retaliate against you for filing a complaint.

We are required by law to maintain the privacy of, and provide individuals with, this notice of our legal duties and privacy practices with respect to protected health information. If you have any objections to this form, please ask to speak with our HIPAA Compliance Officer in person or by phone at our Main Phone Number.

Signature below is only acknowledgement that you	have received this Notice of our Privacy Practices:
Signature	Date

PLEASI NAMI OCCU	E FILL (E: PATIO	OUT THIS FORM O	COMPLETELY SO V AGE:	VE CAN HELP YOU WITH SEX: M / F HEIG DATE LAST V	H YOUR PAINFUL CONDI GHT'" WEIGHT WORKED//	TION. lbs
<u>HPI:</u>				<u>NG</u> IN THE AFFECTED AF LL APPLICABLE DESCRIP	REAS AND <u>PLACING AN 'X</u> TIVE WORDS.	ON THE
CONS	TANT	TEMPORARY	R	P PI		
D AILY	7	OCCASIONAL	一个的出	W WAL		/
A CHIN	1G	SHOOTING		·10 (p 1 //)·	M }	
BURNI	ING	S TABBING	利川州	1 162 61 120 17		7
N UMB	ING	TINGLING		ΛΛ }Λ/ H	={ (
PINS/N	VEEDLE	S			$\langle \ \ \rangle$	3
		-		ING PERCENTAGES. YO % RIGHT LEG PAIN = 100	OUR TOTAL PAIN SHOULD	D ADD UP T
HEAD_	%	Neck	%	RIGHT ARM%	LEFT ARM	<u>%</u>
Снест	%	A BDOME	N%	UPPER BACK%	MID BACK	<u></u> %
Lower	R B ACK	% H IPS/ B U	TTOCKS%	RIGHT LEG%	Left Leg	%
CIRCLE	THE A	CTIVITIES WHICH	TEND TO <u>INCREA</u>	<u>SE</u> YOUR PAIN:		
W ALKI	ING	LIFTING B	ENDING TWIS	STING STANDING	SITTING	
FILL IN	ACTIV	ITIES WHICH <u>DEC</u>	<u>'REASE</u> YOUR PAIN	I:		
OOES T	HIS PAI	N AFFECT YOUR S	SLEEP? YES/NO			
CURRE	NT <i>PAI</i> N	MEDICATIONS:_			PRESCRIBED BY DR.	
MARK	YOUR <u>A</u>	VERAGE PAIN SC	ORE:	-68	010	
		<i>VORST</i> PAIN SCOP		.,	10	
	-1	23	45	68		
			· 	YOUR <u>WORST</u> IMAGINA	·	
		S PAIN BEGIN? IBE HOW IT BEGA		EOUS/ INJURY/ ACCIDE	ENT/ SURGERY/ OTHER)	

	TYPES OF TREATMENT HAVE CIRCLE ALL THAT APPLY.	E YOU HAD IN THE PA	AST TO TREAT YOUR CURRENT PAIN.
PAIN CLINIC PHYSI	CAL THERAPY MASSAGE	CHIROPRACTIC	INJECTIONS SURGERY
ACUPUNCTURE MA	GNETS HERBS OTHER:		
PLEASE <u>LIST</u> ALL TRE	ATMENT BELOW:		
HEALTH CARE PROFESSIONAL			TREATMENT/ MEDICATIONS
DIAGNOSTIC EXAMINA	ATIONS: PLEASE CIRCL	E ALL THAT APPLY	
XRAY CT-SCAN MRI	Myelogram Emg/Ncv O	THER	
PAST/CURRENT MEDIC	AL HISTORY: PLEASI	TIRITH ALL THAT A	
	BLEEDING PROBLEMS CA	NCER CIRRHOSIS	COLITIS DIABETES EMPHYSEMA
HEART TROUBLE H	BLEEDING PROBLEMS CA	NCER CIRRHOSIS (COLITIS DIABETES EMPHYSEMA E HIV/AIDS KIDNEY DISEASE
HEART TROUBLE HI	BLEEDING PROBLEMS CAR EPATITIS / JAUNDICE HIG STROKE THYROID T	NCER CIRRHOSIS (GH BLOOD PRESSURI ROUBLE URINATING	COLITIS DIABETES EMPHYSEMA E HIV/AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE
HEART TROUBLE HI MURMUR SEIZURE ARE YOU TAKING ANY I	BLEEDING PROBLEMS CAR EPATITIS / JAUNDICE HIG STROKE THYROID T BLOOD THINNERS SUCH AS	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HI MURMUR SEIZURE ARE YOU TAKING ANY I	BLEEDING PROBLEMS CAR EPATITIS / JAUNDICE HIG STROKE THYROID T	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HI MURMUR SEIZURE ARE YOU TAKING ANY I LIST ALL CURRENT MED	BLEEDING PROBLEMS CAR EPATITIS / JAUNDICE HIG STROKE THYROID T BLOOD THINNERS SUCH AS	NCER CIRRHOSIS GH BLOOD PRESSURI ROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY HELE LIST ALL CURRENT MED ALLERGIES TO MEDICA	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIG STROKE THYROID T BLOOD THINNERS SUCH AS A PICATIONS: FIONS:	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY HELE ALL CURRENT MEDICATED TO M	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIG STROKE THYROID T BLOOD THINNERS SUCH AS A PICATIONS: FIONS:	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IN LIST ALL CURRENT MEDITALITY ALLERGIES TO MEDICAL PAST SURGICAL HISTOR	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID T BLOOD THINNERS SUCH AS A FICATIONS: FIONS: RY: (PLEASE LIST ALL OPER	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HEART TROUBLE HEART TROUBLE HEART MURMUR SEIZURE ARE YOU TAKING ANY HEART MEDICATE TO MEDICATE TO MEDICATE TO MEDICATE HEART SURGICAL HISTORY SH: SINGLE MARRIE	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID TO BLOOD THINNERS SUCH AS CONTROL OF THE PROBLEM	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFA	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IS LIST ALL CURRENT MED ALLERGIES TO MEDICATE PAST SURGICAL HISTOR CH: SINGLE MARRIE PACKS OF CIGARETTES	BLEEDING PROBLEMS CAR EPATITIS / JAUNDICE HIG STROKE THYROID T BLOOD THINNERS SUCH AS A PICATIONS: EIONS: EY: (PLEASE LIST ALL OPER ED DIVORCED SEP SMOKED/DAY	NCER CIRRHOSIS OF ALCOHO THE PROPERTY OF THE PROPERTY	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO HAD). WED # OF CHILDREN
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IS LIST ALL CURRENT MEDICATE ALLERGIES TO MEDICATE PAST SURGICAL HISTOR CH.: SINGLE MARRIE PACKS OF CIGARETTES SHISTORY OF SUBSTANCE	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID TO BLOOD THINNERS SUCH AS OPERATIONS: FIONS: EY: (PLEASE LIST ALL OPERATIONS) EMANDED DIVORCED SEP SMOKED/DAY ABUSE YES / NO IF YES, W	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFALTIONS YOU HAVE HEARATED WIDOW # OF ALCOHOLD HAT TYPE?	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO HAD). WED # OF CHILDREN OLIC BEVERAGES/DAY
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IS LIST ALL CURRENT MED ALLERGIES TO MEDICATE PAST SURGICAL HISTOR PACKS OF CIGARETTES SHISTORY OF SUBSTANCE THE LIST ANY ILLNESSE	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID TO BLOOD THINNERS SUCH AS EVALUATIONS: FIONS: EY: (PLEASE LIST ALL OPER ED DIVORCED SEP SMOKED/DAY E ABUSE YES / NO IF YES, WES WHICH RUN IN YOUR FAME	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFALTONS YOU HAVE HAT TYPE?	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO HAD)
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IS LIST ALL CURRENT MEDITAL CURRENT MEDITAL PAST SURGICAL HISTOR CH: SINGLE MARRIED MARRIE	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID TO BLOOD THINNERS SUCH AS EVALUATIONS: EVALUATIONS	NCER CIRRHOSIS OF BLOOD PRESSURING TROUBLE URINATING COUMADIN, WARFALTONS YOU HAVE HAT TYPE?	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO HAD) WED # OF CHILDREN OLIC BEVERAGES/DAY
HEART TROUBLE HIS MURMUR SEIZURE ARE YOU TAKING ANY IS LIST ALL CURRENT MEDICAL PAST SURGICAL HISTOR CH: SINGLE MARRIE CACKS OF CIGARETTES SHISTORY OF SUBSTANCE CH: LIST ANY ILLNESSES COS: PLEASE CIRCLE ALL COUGH DIARRHEA / CO	BLEEDING PROBLEMS CARE EPATITIS / JAUNDICE HIGH STROKE THYROID TO BLOOD THINNERS SUCH AS EPATITIS / JAUNDICE HIGH BLOOD THINNERS SUCH AS EPATITIONS: EXTENDED DIVORCED SEPATION SEPATION SEPATION OF YES, WE WHICH RUN IN YOUR FAMILE SYMPTOMS YOU MAY CUIT CONSTIPATION DIZZINESS	NCER CIRRHOSIS OF BLOOD PRESSURING COUMADIN, WARFAL COUMADIN, WARFAL ATIONS YOU HAVE HAT TYPE? LY: REENTLY HAVE: EASY BLEEDING	COLITIS DIABETES EMPHYSEMA E HIV / AIDS KIDNEY DISEASE G ULCER VASCULAR DISEASE RIN, PLAVIX, OR TICLID? YES / NO HAD) WED # OF CHILDREN OLIC BEVERAGES/DAY CHANGE IN VISION CHEST PAIN

Diplomate of American Board of Anesthesiology Practice of Anesthesiology Practice of Anesthesiology and Pain Medicine CONTROLLED SUBSTANCE PATIENT AGREEMENT FORM Wewant to ensure that patients and caregivers have clear communication and safe, effective procedures when patient opioids. For the purposes of this document "I" will be synonymous with the patient receiving one or more entrolle substances, and "provider" will be considered synonymous with the prescriber of these medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most petients do not have serious side effects or drug interactions. Unfortunately, some do experience effects and must stop the medication(s). Common side effects include constipation, fiching, nausea, vomiting, seddion o lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breating (sepecially froy have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreas testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is clopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and crawing, and continued use despite harm or risk to the person. When peop	Diplomate of American Board of Anesthesiology Practice of Anesthesiology and Pain Medicine CONTROLLED SUBSTANCE PATIENT AGREEMENT FORM Wewant to ensure that patients and caregivers have clear communication and safe, effective procedures when patien opioids. For the purposes of this document "I" will be synonymous with the prescriber of these medications. EFFECTIVENESS: For most patients and pain conditions, opioids are effective pain-relieving medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience of effects and must stop the medication(s). Common side effects include constipation, itching, nausea, vorniting, selation or lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing ceptainly by have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrea testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn dild dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdomial and even scieures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a ser percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of dwg us compulsive use and crawing, and continued use despite harm or risk to the person. When people are addicted, they are not t	me	-
We want to ensure that patients and caregivers have clear communication and safe, effective procedures when patient opioids. For the purposes of this document "!" will be synonymous with the patient receiving one or more controlle substances, and "provider" will be considered synonymous with the prescriber of these medications. EFFECTIVENESS: For most patients and pain conditions, opioids are effective pain-relieving medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience seffects and must stop the medication(s). Common side effects include constipation, itching, nausea, vomiting, sedation o lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especialy if yor have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, desiring of sex drive, decreas testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply	Wewant to ensure that patients and caregivers have clear communication and safe, effective procedures when patien opioids. For the purposes of this document "" will be synonymous with the patient receiving one or more controlle substances, and "provider" will be considered synonymous with the prescriber of these medications. EFFECTIVENESS: For most patients and pain conditions, opioids are effective pain-relieving medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience is effects and must stop the medication(s). Common side effects include constipation, fitching, nausea, vorniting, sedation of lighthreadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yo have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrea testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to	e	Diplomate of American Board of Anesthesiology
opioids. For the purposes of this document "i" will be synonymous with the patient receiving one or more controlle substances, and "provider" will be considered synonymous with the prescriber of these medications. EFFECTIVENESS: For most patients and pain conditions, opioids are effective pain-relieving medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTIS: Most patients do not have serious side effects include constipation, itching, nausea, vomiting, sedation o lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yor have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreast testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal, and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug ust compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life. Lower your pain. TREATMENT OPTIONS: Medications.	opioids. For the purposes of this document "I" will be synonymous with the patient receiving one or more controlle substances, and "provider" will be considered synonymous with the prescriber of these medications. EFFECTIVENESS: For most patients and pain conditions, opioids are effective pain-relieving medications. However, it possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience is effects and must stop the medication(s). Common side effects include constipation, itching, nausea, vomiting, sedation ol lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yo have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrea testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your pain. TREATMENT OPTIONS: Reali	CONTROLLED SUBSTANC	CE PATIENT AGREEMENT FORM
possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience seffects and must stop the medication(s). Common side effects include constipation, tiching, nausea, vomiting, seation or lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially fy or have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreast testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO:	possible opioids will not work well for you and your pain. SAFETY: Most people can take these drugs safely, but some people do experience side effects. (See below.) SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience effects and must stop the medication(s). Common side effects include constipation, itching, nausea, vomiting, sedation or lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yo have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreat testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Counseling relaxation training, hypnosis and meditation, Remain active every da	opioids. For the purposes of this	s document "I" will be synonymous with the patient receiving one or more controlled
SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience seffects and must stop the medication(s). Common side effects include constipation, itching, nausea, vomiting, sedation of lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if you have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrease testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdeminally and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	SIDE EFFECTS: Most patients do not have serious side effects or drug interactions. Unfortunately, some do experience effects and must stop the medication(s). Common side effects include constipation, itching, nausea, vomiting, sedation or lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yo have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrea testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medication		
effects and must stop the medication(s). Commonside effects include constipation, itching, nausea, vomiting, sedation of lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if you have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreast estosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal yand even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	effects and must stop the medication(s). Commonside effects include constipation, itching, nausea, vomiting, sedation or lightheadedness. Uncommon reactions include swelling in the legs, water on the lungs, trouble breathing (especially if yo have emphysema/COPD or are on other narcotics), mental slowing and loss of coordination, lowering of sex drive, decrea testosterone (male sex hormone) and addiction. Note: Pregnant women using opioids could make their newborn child dependent upon opioids. If you are pregnant, you need to alert your health care provider. DEPENDENCE: Dependence is not the same as addiction. Many people who take opioids daily will become dependent on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schi Fini	SAFETY: Most people can take the	ese drugs safely, but some people do experience side effects. (See below.)
on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication is stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	on them. Dependence is when your body adapts to the medication and then experiences withdrawal if the medication stopped or lowered too quickly. Withdrawal symptoms include moodiness, aches and pains, sweating, diarrhea, abdominal and even seizures. ADDICTION: Addiction is not the same as dependence. While many people become dependent on daily opioids, only a sm percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug us compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. The provider of the provider and staff have explained the risks and benefits of chronic opioid therapy for my paints.	effects and must stop the medicat lightheadedness. Uncommon read have emphysema/COPD or are on testosterone (male sex hormone) a	tion(s). Common side effects include constipation, itching, nausea, vomiting, sedation or ctions include swelling in the legs, water on the lungs, trouble breathing (especially if you other narcotics), mental slowing and loss of coordination, lowering of sex drive, decreased and addiction. Note: Pregnant women using opioids could make their newborn child
percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life. Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	percentage of these people will become addicted. Addiction is characterized by behaviors such as loss of control of drug use compulsive use and craving, and continued use despite harm or risk to the person. When people are addicted, they are not taking opioids simply to treat the pain. GOALS: The goals of chronic pain management are to: Improve your ability to function in your daily life, Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. The Schifini and staff have explained the risks and benefits of chronic opioid therapy for my pain.	on them. Dependence is when you stopped or lowered too quickly. With	ur body adapts to the medication and then experiences withdrawal if the medication is
 Improve your ability to function in your daily life, Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. 	 Improve your ability to function in your daily life. Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini 	percentage of these people will beco	ome addicted. Addiction is characterized by behaviors such as loss of control of drug use, intinued use despite harm or risk to the person. When people are addicted, they are not
 2 Lower your pain. TREATMENT OPTIONS: Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. 	 2 Lower your pain. TREATMENT OPTIONS: Medications, 2 Counseling, relaxation training, hypnosis and meditation, 3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: 1 Realize that opioid therapy is only one part of treatment. 2 Remain active every day and try to increase activity a little bit at a time. 3 Use your medications ONLY as directed by your provider. 4 Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	GOALS: The goals of chronic pain m	nanagement are to:
TREATMENT OPTIONS: 1 Medications, 2 Counseling, relaxation training, hypnosis and meditation, 3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: 1 Realize that opioid therapy is only one part of treatment.	TREATMENT OPTIONS: ① Medications, ② Counseling, relaxation training, hypnosis and meditation, ③ Chiropractic care, massage, acupuncture and physical therapy, ④ Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: ① Realize that opioid therapy is only one part of treatment. ② Remain active every day and try to increase activity a little bit at a time. ③ Use your medications ONLY as directed by your provider. ④ Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page	1 Improve your ability to function in	n your daily life,
 1 Medications, 2 Counseling, relaxation training, hypnosis and meditation, 3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: 1 Realize that opioid therapy is only one part of treatment. 	 Medications, Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	2 Lower your pain.	
 2 Counseling, relaxation training, hypnosis and meditation, 3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. 	 Counseling, relaxation training, hypnosis and meditation, Chiropractic care, massage, acupuncture and physical therapy, Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Or. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	TREATMENT OPTIONS:	
3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	 3 Chiropractic care, massage, acupuncture and physical therapy, 4 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: 1 Realize that opioid therapy is only one part of treatment. 2 Remain active every day and try to increase activity a little bit at a time. 3 Use your medications ONLY as directed by your provider. 4 Work with your provider and follow treatment recommendations in addition to taking prescribed medications. 2 Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	Medications,	
Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	 Surgery and spinal or other type of injections. WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	2 Counseling, relaxation training, h	ypnosis and meditation,
WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment.	WHAT YOU NEED TO DO: Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. or. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page.	3 Chiropractic care, massage, acup	puncture and physical therapy,
Realize that opioid therapy is only one part of treatment.	Realize that opioid therapy is only one part of treatment. Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page.	4 Surgery and spinal or other type	of injections.
	Remain active every day and try to increase activity a little bit at a time. Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page.	WHAT YOU NEED TO DO:	•
Remain active every day and try to increase activity a little bit at a time.	 Use your medications ONLY as directed by your provider. Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page. 	Realize that opioid therapy is on	nly one part of treatment.
_	Work with your provider and follow treatment recommendations in addition to taking prescribed medications. Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my page.	Remain active every day and try t	o increase activity a little bit at a time.
3 Use your medications ONLY as directed by your provider.	Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my pa	3 Use your medications ONLY as di	irected by your provider.
Work with your provider and follow treatment recommendations in addition to taking prescribed medications.		Work with your provider and follows:	ow treatment recommendations in addition to taking prescribed medications.
Dr. Schifini and staff have explained the risks and benefits of chronic opioid therapy for my pa	I,, understand that I must comply with the following rules or I will not be given opioi	Dr. Schifini	_and staff have explained the risks and benefits of chronic opioid therapy for my pain.
I,, understand that I must comply with the following rules or I will not be given opioid		l,	, understand that I must comply with the following rules or I will not be given opioids.

continued on page 2 of 2 >-

I will take the medication(s) listed in my charts as they were prescribed and intended and only in that manner.

Name	
Date	Joseph J. Schifini, M.D., Ltd Diplomate of American Board of Anesthesiology Practice of Anesthesiology and Pain Medicine
*Initial in each box	below.
	se the dose or stop the medication unless asked to do so by my provider or my provider's worrisome side effect soon after it begins.
physical and occupation	ough on appointments that may help me with chronic pain and functioning. These may include all therapy, counseling and other mental health practices, neurosurgery, neurology and orthopedics. p these appointments and therapies may result in the stopping of the opioid medications.
lf prescribed, I w	vill use medications other than opioids to control pain.
I will accept opic	oids for chronic pain from my provider only.
	exchange or sell my opioids, as the law prohibits those actions. I understand that my provider as of drug misuse to any and all authorities for investigation.
I will not use illega provider.	al/street drugs (this includes marijuana). I will not use narcotic medications unless provided to mefrom my
l agree to provide	samples for random drug testing when asked. If I fail to provide the sample when asked or if the I may forfeit the right to continue receiving the medication.
	concerned that I might have a substance abuse problem, I must agree to an evaluation by a on. If the evaluation suggests I have a drug abuse problem, my provider may stop my medication in withdrawal symptoms.
I will not get early re	efills unless something has dramatically changed and then only if my provider agrees.
	oioids by themselves, in combination with alcohol or in combination with other medicationscan nd loss of coordination. I agree to contact my provider if these symptoms arise. I should not if I have these side effects.
not be refilled early. Each c	lity to keep my medications safe. If opioids are lost, damaged or stolen, the medication may or may ase will be looked at individually. If the medication is stolen, I must file a police report and submit to my provider's office. Again, stolen medications may or may not be refilled. If a refill is given, it
	develops that causes acute pain, I have the right to expect appropriate treatment for that new treating me for the new condition. I should not be required to increase the use of my chronic s and new pain.
I understand that if may lower or stop the opioid	my provider does not feel I am following through adequately with the treatment plan, my provider altogether.
I understand that m have not responded positivel	y provider may decide to stop the opioid if after increasing it adequately, my pain and function ly.
and involved with the invest past caregivers, all pharmac	orize my provider's office to contact any and all groups and organizations involved with my care igation of medication and drug abuse. I give permission to my provider to discuss my care with sies and policing agencies. This also gives these caregivers and pharmacies permission to share about my past treatments and care.
PATIENT SIGNATURE	DATE
HEALTH CARE PROVIDER	DATE

Patient's Name

Controlled Sub	ostance Questionnaire		<u>YES</u>	<u>NO</u>	N/A
N/A means not applical	ble.				
Have you ever used a con	trolled substance in a way other than prescri	bed?			
Have you ever diverted a	controlled substance to another person?				
Have you ever taken a cor	strolled substance that did not have the desir	ed effect?			
Are you currently using an	y drugs, including alcohol or marijuana?				
Are you using any drugs th	at may negatively interact with a controlled	substance?			***********
Are you using any drugs th	at were not prescribed by a practitioner that	is treating you?			•
Have you ever attempted t	o obtain an early refill of a controlled substa	nce?			
Have you ever made a clair	m that a controlled substance was lost or sto	len?			
Have you ever been questi	oned about your pharmacy report or PMP re	port?			
Have you ever had blood o	r urine tests that indicate inappropriate usag	e of meds?		-	
Have you ever been accuse	d of inappropriate behavior or intoxication?				***********
Have you ever increased th	e dose or frequency of meds without telling	your provider?		************	-
Have you ever had difficulty	with stopping the use of a controlled substa	ance?		******	
Have you ever demanded to	be prescribed a controlled substance?			*****************	-
Have you ever refused to co	operate with any medical testing or examina	ations?	MARKET PARTY OF THE PARTY OF TH		
Have you ever had a history	of substance abuse of any kind?				
Has there been any change	in your health that might affect your medicat	tions?			
Have you misused or becom	e addicted to a drug, or failed to comply with	h instructions?			
Are there any other factors t	that your practitioner should consider before	e prescribing?	•	to a second distribution of the second of th	***********
		Manage And Adjusted States			
Patient's Signature	Patient's Printed Name	Date			
Parent/Legal Guardian	Parent/Legal Guardian	 Date			

PRESCRIPTION OPIOID MISUSE INDEX

1. Do you ever use MORE of your medication, that is, tal prescribed for you?	ke a hi Yes	_	sage, th	an is
2. Do you ever use your medication MORE OFTEN, that dosages?	is, sho		e time b	etween
3. Do you ever feel high or get a buzz after using your pai	in med	lication?	Yes	No
4. Do you ever take your pain medication because you are relieve or cope with problems other than pain?	upset Yes	, using t No	he medi	cation to
5. Have you ever gone to multiple physicians including emmore of your pain medication?	nergen Yes	•	doctors	, seeking
6. Do you ever need early refills for your pain medication?	Yes	s No		
Name				
Date				

Name		
Date		Poolele Depression Inventory
	This d	Beck's Depression Inventory epression inventory can be self-scored. The scoring scale is at the end of the questionnaire.
	· _ I.	oprocessor inventory sair be seri-secred. The secret is at the end of the questionname.
	~ 7\$	0 I do not feel sad.
		1 I feel sad
		I am sad all the time and I can't snap out of it.
		I am so sad and unhappy that I can't stand it.
	2.	
		I am not particularly discouraged about the future.
\		I feel discouraged about the future.
		2 I feel I have nothing to look forward to.
		I feel the future is hopeless and that things cannot improve.
	3.	
		I do not feel like a failure.
		I feel I have failed more than the average person.
		As I look back on my life, all I can see is a lot of failures.
		I feel I am a complete failure as a person.
	4.	
		I get as much satisfaction out of things as I used to.
	2	- ,
	3	I am dissatisfied or bored with everything.
	5.	I doubt fool moution house miles
	0 1	I don't feel particularly guilty I feel guilty a good part of the time.
	2	I feel quite guilty most of the time.
	3	I feel guilty all of the time.
	6.	Theorem and of the time.
	0.	I don't feel I am being punished.
	1	I feel I may be punished.
	2	I expect to be punished.
	. 3	I feel I am being punished.
	7.	O F
	0	I don't feel disappointed in myself.
	1	I am disappointed in myself.
	2	I am disgusted with myself.
	3	I hate myself.
	8.	
	0	I don't feel I am any worse than anybody else.
	1	I am critical of myself for my weaknesses or mistakes.
	2	I blame myself all the time for my faults.
	3	I blame myself for everything bad that happens.
	9.	
	0	I don't have any thoughts of killing myself.
	1	I have thoughts of killing myself, but I would not carry them out.
	2	I would like to kill myself.
	3	I would kill myself if I had the chance.
	10.	
	0	I don't cry any more than usual.
	1	I cry more now than I used to.
	2	I cry all the time now.
	3	I used to be able to cry, but now I can't cry even though I want to.

Name	
Date	
. 11.	
0	I am no more irritated by things than I ever was.
1	I am slightly more irritated now than usual.
2	I am quite annoyed or irritated a good deal of the time.
3 12.	I feel irritated all the time.
12.	I have not lost interest in other people.
1	I am less interested in other people than I used to be.
0	I have lost most of my interest in other people.
\ 2 3	I have lost all of my interest in other people.
13.	
0	I make decisions about as well as I ever could.
1	I put off making decisions more than I used to.
2	I have greater difficulty in making decisions more than I used to.
3	I can't make decisions at all anymore.
14.	
0	I don't feel that I look any worse than I used to.
. 1	I am worried that I am looking old or unattractive.
. 2	I feel there are permanent changes in my appearance that make me look
3	unattractive
15.	I believe that I look ugly.
. 0	I can work about as well as before.
1	It takes an extra effort to get started at doing something.
2	I have to push myself very hard to do anything.
3	I can't do any work at all.
16.	•
0	I can sleep as well as usual.
1	I don't sleep as well as I used to.
2	I wake up 1-2 hours earlier than usual and find it hard to get back to sleep.
3	I wake up several hours earlier than I used to and cannot get back to sleep.
17.	
0	I don't get more tired than usual.
Ī	I get tired more easily than I used to.
2	I get tired from doing almost anything.
3	I am too tired to do anything.
18.	, 0
0	My appetite is no worse than usual.
1	My appetite is not as good as it used to be.
2	My appetite is much worse now.
3	I have no appetite at all anymore.
19.	
0	I haven't lost much weight, if any, lately.
1	I have lost more than five pounds.
2 3	I have lost more than ten pounds.
3	I have lost more than fifteen pounds.

wame_			· ·
Date			
		20.	
		- : 0	I am no more worried about my health than usual.
		1	I am worried about physical problems like aches, pains, upset stomach, or constipation.
		2	I am very worried about physical problems and it's hard to think of much else.
		3	I am so worried about my physical problems that I cannot think of anything else.
		21.	
		0	I have not noticed any recent change in my interest in sex.
	\	1	I am less interested in sex than I used to be.
		2	I have almost no interest in sex.
		2	I have lost interest in say completely

INTERPRETING THE BECK DEPRESSION INVENTORY

Now that you have completed the questionnaire, add up the score for each of the twenty-one questions by counting the number to the right of each question you marked. The highest possible total for the whole test would be sixty-three. This would mean you circled number three on all twenty-one questions. Since the lowest possible score for each question is zero, the lowest possible score for the test would be zero. This would mean you circles zero on each question. You can evaluate your depression according to the Table below.

Total Score	Levels of Depression
1-10	These ups and downs are considered normal
11-16	Mild mood disturbance
17-20	Borderline clinical depression
21-30	Moderate depression
31-40	Severe depression
over 40	Extreme depression